

COMMON SLEEPING TABLETS, PAINKILLERS & ANXIETY MEDICATION

What are the best sleeping tablets for treating insomnia? What is the difference between Zopiclone and Zolpidem tablets? Which anxiety medication works the fastest? To help you decide on the best treatment for your insomnia, pain or anxiety see our quick reference guide and compare the most popular [sleeping tablets](#), [painkillers](#) and [anxiety tablets](#) that are [FDA](#) and [EMA](#) approved and available now for speedy home delivery.

NAME / BRAND	TYPE	TREAT	FORM	DOSE	CONSUME	ONSET	DURATION	ADVICE
Zopiclone <i>Imovane / Actavis</i> i	Non-benzodiazepne Sleeping Tablets	Insomnia	Tablet / Pill	3.75mg, 7.5mg, 10mg	1 x Zopiclone Tablet per Day (Before Bed)	40 - 60 min	8 - 9 hours	Zopiclone tablets may be consumed with or without food.
Zolpidem Tartrate <i>Ambien / Ambien CR</i> i	Non-benzodiazepine Sleeping Tablets	Insomnia	Tablet / Pill	5mg, 10mg	1 x Zolpidem Tablet per Day (Before Bed)	30 - 40 min	8 - 9 hours	Zolpidem tablets may be consumed with or without food.
Temazepam <i>Restoril / Teva</i> i	Benzodiazepine Sleeping Tablets	Insomnia	Tablet / Capsule	10mg, 20mg, 30mg	1 x Temazepam Tablet per Day (Before Bed)	30 - 40 min	7 - 8 hours	Temazepam tablets may be consumed with or without food.
Nitrazepam <i>Mogadon / Actavis</i> i	Benzodiazepine Sleeping Tablets / Anxiety Aid	Insomnia & Anxiety	Tablet / Pill	5mg, 10mg	1 x Nitrazepam Tablet per Day as Required	30 - 60 min	6 - 8 hours	Nitrazepam tablets may be consumed with or without food.
Diazepam <i>Valium / Bensedin</i> i	Benzodiazepine Sleeping Tablets / Anxiety Aid	Insomnia & Anxiety	Tablet / Pill	2mg, 5mg, 10mg	1 x Diazepam Tablet 2 - 4 Times per Day as Required	40 - 60 min	4 - 6 hours	Diazepam tablets may be consumed with or without food.
Alprazolam <i>Xanax / Pfizer</i> i	Benzodiazepine Anxiety Aid	Anxiety & Panic Attacks	Tablet / Pill	0.5mg, 1mg, 2mg	1 x Alprazolam Tablet 2 - 4 Times per Day as Required	40 - 60 min	4 - 5 hours	Alprazolam tablets may be consumed with or without food.
Lorazepam <i>Ativan / Pfizer</i> i	Benzodiazepine Anxiety Aid	Anxiety & Panic Attacks	Tablet / Pill	0.5mg, 1mg, 2.5mg	1 x Lorazepam 2 - 3 Times per Day as Required	20 - 30 min	6 - 8 hours	Lorazepam tablets may be consumed with or without food.
Clonazepam <i>Klonopin / Rivotril</i> i	Benzodiazepine Anxiety Aid	Anxiety & Panic Attacks	Tablet / Pill	1mg, 2mg	1 x Clonazepam Tablet 1 - 2 Times per Day as Required	20 - 60 min	3 - 4 hours	Clonazepam tablets may be consumed with or without food.
Tramadol <i>Ultram / Mylan</i> i	Opiate Analgesic Painkiller	Moderate to Severe Pain	Tablet / Pill	50mg, 100mg	1 x Tramadol Tablet 2 - 4 Times per Day as Required	30 - 60 min	4 - 6 hours	Tramadol tablets may be consumed with or without food.
Dihydrocodeine <i>Actavis / Almus</i> i	Analgesic Combination Painkiller	Moderate to Severe Pain	Tablet / Pill	16mg/ 356.4 mg/30mg	1 x Dihydrocodeine Tablet 2 - 3 Times per Day as Required	60 - 90 min	4 - 6 hours	Dihydrocodeine tablets may be consumed with or without food.
Codeine Phosphate <i>Teva / Wockhardt</i> i	Opiate Analgesic Painkiller	Mild to Moderate Pain	Tablet / Pill	30mg, 60mg	2 - 4 Times per Day as Required	30 - 60 min	4 - 6 hours	Codeine Phosphate tablets may be consumed with or without food.